

## BAB 6. KESIMPULAN DAN SARAN

### 6.1. Kesimpulan

Hasil penelitian ini dapat disimpulkan bahwa substitusi butiran kering destilat (BKD) produk ikutan produksi bioetanol oleh ko-kultur *S. cerevisiae* dengan *C. tropicalis* dari sekam padi dalam formula pakan puyuh dapat mempengaruhi perubahan gas ammonia dalam kandang dan kadar air feses. Pemberian pakan puyuh dengan substitusi 10% BKD sekam padi dapat mengurangi kadar gas ammonia berbahaya dalam kandang unggas sebesar 3.505% dan kandungan air feses 10% lebih rendah. Substitusi butiran kering destilat (BKD) produk ikutan produksi bioetanol oleh ko-kultur *S. cerevisiae* dengan *C. tropicalis* dari sekam padi dalam formula pakan puyuh juga dapat mempengaruhi kandungan unsur karbon (C), nitrogen (N), kalsium (Ca) dan phosphor (P) pada feses puyuh. Substitusi 20% BKD sekam padi dalam formulasi pakan puyuh dapat meningkatkan penyerapan unsur fosfor (P) di usus. Substitusi 10% BKD sekam dalam formulasi pakan puyuh tidak mempengaruhi penyerapan karbon (C), meningkatkan penyerapan fosfor (P) tetapi menurunkan penyerapan nitrogen (N) dan sedikit kalsium (Ca).

### 6.2. Saran

Berdasarkan kesimpulan dalam penelitian ini, untuk meningkatkan kesehatan lingkungan dan meningkatkan penyerapan nutrisi dalam usus puyuh disarankan menggunakan butiran kering destilat (BKD) dari sekam padi sebesar 10% yang disubstitusi dalam ransum puyuh yang sedang bertelur.

## UCAPAN TERIMA KASIH

Peneliti mengucapkan terima kasih kepada Rektor Universitas 17 Agustus 1945 Surabaya melalui LPPM Untag Surabaya yang telah membiayai kegiatan Penelitian dengan judul “Penurunan Emisi Gas Dalam Kandang Puyuh dan Karakteristik Kimia Feses Unggas Yang Diberi Butiran Kering Destilat Produk Ikutan Produksi Bioetanol Oleh Ko-Kultur *S. cerevisiae* Dengan *C. tropicalis* dari Sekam Padi” Tahun Anggaran 2019 dengan Kontrak Penugasan No. 487.39/ST/003/LPPM/ Lit/VII/2019. Peneliti juga mengucapkan terima kasih kepada kepala Lab. Nutrisi FKH-UA dan Environmental Laboratory, Mechanical Laboratory

and Calibration Mutiara Kebonagung yang telah memberikan kesempatan dan waktunya dalam melakukan analisis bahan penelitian.

## DAFTAR PUSTAKA

- Agarwal, N., C. Shekhar, R. Kumar, L. C. Chaudhary and D. N. Kamra. (2009) Effect of peppermint (*Mentha piperita*) oil on in vitro methanogenesis and fermentation of feed with buffalo rumen liquor. *Anim. Feed Sci. Technol.* 148:321-327.
- Abousekken. M.S.M. 2014. Use of corn distillers dried grains with solubles (DDGS) inlaying quail diets. *Egypt. Poult. Sci.* Vol. 34 (3): 681-703. <http://www.epsaegypt.com>.
- Abraham, E., Deepa, B., L. A., Jacob, M., Thomas, S., Cvelbar, U., et al., 2011. Extraction of nanocellulose fibrils from lignocellulosic fibres: a novel approach. *Carbohydrate Polymers*, 86, 1468-1475.
- Agus, A. 2007. Membuat Pakan Ternak Secara Mandiri. PT Aji Parama, Yogyakarta.
- Amerah, A. M., V. Ravindran, R. G., Lentle and D. G. Thomas. 2007. Feed particle size : implication on the digestion and performance of poultry. *J. World's Poultry. Sci.* 63: 439-453.
- Amrullah, I. K. 2003. Nutrisi Ayam Petelur. Lembaga Satu Gunung Budi, Bogor.
- Anggorodi, R. 1985. Ilmu Makanan Ternak Unggas : Kemajuan Mutakhir. Universitas Indonesia Press, Jakarta.
- Anggorodi, R. 1994. Ilmu Makanan Ternak Umum. Penerbit PT. Gramedia, Jakarta
- Applegate, T.J., C. Troche, ., Z. Jiang, Z. and T. Johnson. 2009. The nutritional value of high protein corn distillers dried grains for broiler chickens and its effect on nutrientexcretion. *Poul. Sci.* 88: 354-359.
- Batal, A. B. and N. M. Dale. 2006. True metabolizable energy and amino acid digestibility of distillers dried grains with solubles. *J. Appl. Poult. Res.* 15:89-93.
- Cosgrove DJ. 1980. Relaxation in a High-Stress Environment. The Molecular Bases of Extensible Cell Walls and Cell Enlargegment. *Plant Cell*. 9: 1031-1041.
- Deniz, G., H. Gencoglu, S.S. Gezen, I.I. Turkmen, A. Orman and C. Kara, 2013. Effects offeeding corn distiller's dried grains with solubles with and without enzyme cocktail supplementation to laying hens on performance, egg quality, selected manure parameters and feed cost. *Livestock Sci.*, 152: 174-181.
- Ezzat M. Abd El-Hack, M. Alagawany, M. R. Farag and K. Dhama. 2015. Use of maize distiller's dried grains with solubles (DDGS) in Laying Hen Diets: Trends and Advances. *Asian J. Anim. Vet. Adv.*, 10 (11): 690-707.
- Fastinger, N.D., J.D. Latshaw, and D.C. Mahan. 2006. Amino Acid Availability and True Metabolizable Energy Content of Corn Distillers Dried Grains with Solubles in AdultCecectomizedRoosters. *Poul. Sci.* 85: 1212-1216.
- Francis, G., Z. Kerem, H. P. S. Makkar and K. Becker. (2002) The biological action of saponin in animal system: A review. *Br. J. Nutr.* 88:587-605.
- Goel, G., H. P. S. Makkar and K. Becker. (2008) Effects of Sesbania sesban and Carduus pycnocephalus leaves and Fenugreek (*Trigonella foenum-graecum L.*) seeds and their extracts on partitioning of nutrients from roughage- and concentrate-based feeds to methane. *Anim. Feed Sci. Technol.* 147:72-89.
- Hassan, S. M. and Al Aqil, A. A. 2015. Effect of adding different dietary levels of distillers dried grains with solubles (DDGS) on productive performance of broiler chicks. *International Journal of Poultry Science* 14 (1), 13-18.
- Hintono, A. 1997. Kualitas Telur yang disimpan dalam Kemasan Atmosfer Termodifikasi. *Jurnal sainteks*. Edisi ke-4. Halaman 45--51.

- Lassey, K. R (2007) Livestock methane emission from the individual grazing animal through national inventories to the global methane cycle. *Agric Forest Meteorol.* 142 : 120-132.
- Loar, R.E. II, J. S. Moritz, J.R. Donaldson, and A. Corzo. 2010. Effects of feeding distillerssoluble to broilers from 0 to 28 days posthatch on broiler performance, feed manufacturing efficiency, andselected intestinal characteristics. *Poul. Sci.* 89:2242-2250.
- Lovett, D. K., L. J. Stack, S. Lovell, J. Callan, B. Flynn, M. Hawkins and F.P. O'Mara. (2005) Manipulating enteric methane emissions and animal performance of late-lactation dairy cows through concentrate supplementation at pasture. *J. Dairy Sci.* 88:2836-2842.
- Makkar, H. P. S. (2005) In vitro gas methods for evaluation of feeds containing phytochemicals. *Anim. Feed. Sci. Technol.* 123:291-302.
- Masa'deh, M.K., S.E. Purdum, and K.J. Hanford. 2011. Dried distillers grains with soluble in laying hen diets. *Poul. Sci.* 90:1960-1966.
- Maynard, L.A. Loosil, J.K. Hintz, H.F and Warner, R.G. 2005 Animal Nutrition. (7th Edition) McGraw-Hill Book Company. New York, USA.
- Moss, A.R., J.P. Jouany and J. Newbold. (2000) Methane production by ruminants: its contribution to global warming. *Ann. Zootech.* 49:231-253.
- Murtidjo, B. A. 2005. Ayam lokal Cetakan ke-5. Kanisius, Yogyakarta.
- Mwenya, B., B. Santoso, C. Sar, Y. Gamo, T. Kobayashi, I. Arai and J. Takahashi. (2004) Effects of including 1–4 galactooligosaccharides, lactic acid bacteria or yeast culture on methanogenesis as well as energy and nitrogen metabolism in sheep. *Anim. Feed Sci. Technol.* 115:313-326.
- National Research Council. (1994) Nutrient Requirements of Poultry. 9th Resived Edition. National Academic Press, Wasington, DC.
- Ning, D., J. M. Yuan, Y. W. Wang, Y. Z. Peng, and Y. M. Guo. 2014. The Net EnergyValues of Corn, Dried Distillers Grains with Solubles and Wheat Bran for LayingHensUsing Indirect Calorimetry Method Asian Australas. *J. Anim. Sci.* Vol. 27, No. 2 : 209-216 February 2014.<http://dx.doi.org/10.5713/ajas.2013.13243>
- North, M.D, and D.D. Bell, 1990.Commercial Chicken Production Manual.Second Edition. The Avi Publishing Co. Inc. Wesport, Conecticut.
- Patiyandela, R. (2013) Kadar NH<sub>3</sub> dan CH<sub>4</sub> Serta CO<sub>2</sub> Dari Peternakan Broiler Pada Kondisi Lingkungan Dan Manajemen Peternakan Berbeda Di Kabupaten Bogor.
- Patra, A. K., D. N. Kamra and N. Agarwal. (2006) Effect of plant extracts on in vitro methanogenesis, enzyme activities and fermentation of feed in rumen liquor of buffalo. *J. Anim. Feed Sci. Technol.* 128:276-291.
- Pauzenga. (1991) Animal P. Inrtoduction in the 90's in harmony with nature, A case study in the Nederlands. In biotechnology in the feed Industry. Proc. Alltech!s Seven Annual Symp. Nicholasville, Kentucky.
- Piliang, W.G., A. Suprayogi, N. Kusmorini, M. Hasanah, S. Yuliani, dan Risfaheri. 2011. Efek Pemberian Daun Katuk (*Sauvopus Androgynus*) dalam Ransum terhadap Kandungan Kolesterol Karkas dan Telur Ayam Lokal. Lembaga Penelitian IPB bekerjasama dengan Badan Penelitian dan Pengembangan Pertanian.
- Purdum, S., Hanford, K. and Kreifels, B. 2014. Short-term effects of lower oil dried distillers grains with solubles in laying hen rations. *Poultry Science* 93(1), 2592-2595.

- Ramírez Restrepo C. A., T. N. Barry, A. Marriner, N. López-Villalobos, E. L. McWilliam, K. R. Lassey and H. Clark. (2010) Effects of grazing willow fodder blocks upon methane production and blood composition in young sheep. *Anim. Feed Sci. Technol.* 155:33-43.
- Ritz, C. W., B. D. Fairchild, & M. P. Lacy. (2004) Implications of ammonia production and emissions from commercial poultry facilities: a review. *J. Appl. Poult. Res.*
- Shim, M.Y., G.M. Pesti, R.I. Bakalli, P.B. Tillman, and R.L. Payne. 2011. Evaluation of DDGS as an alternative ingredient for broiler chickens. *Poult. Sci.* 90:369-376.
- Siregar, A.P. dan M. Sabrani.(1970) Teknik Modern Beternak Ayam. C.V. Yasaguna. Jakarta
- Sopandi, T. dan Wardah, A. 2015. Sugar consumption in mono and co-culture *Saccharomyces cerevisiae* and others selected microorganism for bioethanol production from stream rice husk medium. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences* 17(3), 577-586.
- Sopandi, T. dan Wardah, A. (2017). Ethanol production and sugar consumption of co-culture *Saccharomyces cerevisiae* FNCC 3012 with *Candida tropicalis* FNCC 3033 in media containing inhibitor fermentation. *Journal of Microbiology, Biotechnology and Food Sciences* 7(2), 160-167.
- Standar Nasional Indonesia. (2008) Pakan Ayam Ras Pedaging. SNI 01-3929- 2006.
- Statistik Peternakan dan Kesehatan Hewan, Direktorat Jenderal Peternakan dan Kesehatan Hewan. (2017) Kementerian Pertanian RI. Jakarta
- Steiner, H.H. and G.C. Shurson. 2009. The use and application of distillers dried grains withsolubles in swine diets. *J of Anim. Sci.* 87: 1292-1303.
- Svensson, I. (1990) Putting the lid on the heaps. *Acid. Enviro. Magazine.* 9: 11 – 15
- Sopandi, T dan Wardah. (2019) Production Performance and Carcass Percentage of Broilers Fed Distillers Dried Grain From Rice Husks With Co-culture Fermentation of *Saccharomyces cerevisiae* with *Candida tropicalis*. *International Journal of Poultry Science.* 18 (2) : 80-87.
- Sopandi, T., T. Surtiningsih dan A. Wardah. 2019. Nutrient compositions of distillers dried grain from rice husks with co-culture fermentation of *Saccharomyces cerevisiae* with *Candida tropicalis*. *Malaysian Journal of Microbiology*, Vol 15(3) 2019, pp. 173-181.
- Sudaryani. 2003. Kualitas Telur. Penebar Swadaya. Jakarta.
- Suprijatna, E., U. Atmomarsono, dan R. Kartasudjana. 2005. Ilmu Dasar Ternak Unggas. Penebar Swadaya, Jakarta.
- Tillman, A. D., H. Hartadi, S. Reksohadiprodjo, S. Prawirokusumo dan S. Lebdosoekojo. 1998. Ilmu Makanan Ternak Dasar. Gadjah Mada University Press, Yogyakarta.
- Tillman, A. D. 1991. Komposisi Bahan Makanan Ternak Untuk Indonesia. Gadjah Mada University Press. Yogyakarta.
- Takahashi, J., B. Mwenya, B. Santoso, C. Sar, K. Umetsu, T. Kishimoto, K. Nishizaki, K. Kimura and O. Hamamoto. (2005) Mitigation of methane emission and energy recycling in animal agricultural systems. *Asian Austral. J. Anim. Sci.* 18:1199-1208.
- Ungerfeld, E. M., S. R. Rust, R. J. Burnett, M. T. Yokoyama and J. K. Wang. (2005) Effects of two lipids on in vitro ruminal methane production. *Anim. Feed Sci. Technol.* 119:179-185.
- Wahju, J. 2004. Ilmu Nutrisi Unggas. Cetakan ke lima. Gadjah Mada University Press, Yogyakarta.

- Wang, Z., S. Cerrate, C. Coto, F. Yan and P.W. Waldroup. 2007. Utilization of distillersdried grains with solubles (DDGS) in broiler diets using a standardized nutrient matrix. *Int. J. Poult. Sci.*, 6: 470-477.
- Wardah dan Panjaitan, T. W. S. Panjaitan, (2019) Penurunan Emisi Gas Dalam Kandang Puyuh dan Karakteristik Kimia Feses Unggas Yang Diberi Butiran Kering Destilat Produk Ikutan Produksi Bioetanol Oleh Ko-Kultur *Saccharomyces cerevisiae* dengan *Candida tropicalis* dari Sekam Padi. Hasil Penelitian. Laporan Hasil Penelitian Hibah Perguruan Tinggi. Untag. Surabaya.
- Wardah, Sopandi, T dan Rahmahani, J. (2017) Penggunaan Pakan Fungsional Immunostimulan dan Penurun Kolesterol Telur Berbasis Serbuk Daun Seligi Guna Mengatasi Kendala Ketersedia-an Pakan dan Tingginya Mortalitas pada Puyuh. Laporan Hasil Penelitian Strategis Nasional. Tahun ke 3. Untag. Surabaya.
- Widodo, W. 2002. Peningkatan kualitas bungkil biji karet sebagai bahan pakan ayam pedaging melalui perlakuan fisik dan penambahan kalsium sulfat. Disertasi. Program Pasca Sarjana Universitas Airlangga, Surabaya.
- Youssef, A.W., M.M. El-Moniary and A.H. Abd El-Gawad, 2009. Evaluation of Distiller Dried Grains with soluble (DDGS) as a feedstuff in poultry diets. *Am.-Eurasian J. Agric. Environ. Sci.*, 5: 540-544.